

Synagis Indicator Reports


Last Modified on 03/12/2021 4:05 pm EST

Version 14.19

Overview

If the practice administers Synagis injections, it may be necessary to produce a report of patients for follow-up that have the indicator set. Below are listed several reports that are available. Your practice workflow determines the report that will be used.

Note: For the below reports to produce a patient list, you must set the **Synagis indicated** status on the Indicators tab of the Immunization record.



Forecast Complete List Indicators Notes Forecast Key

Status Indicators by Antigen Influenza / PPV / Synagis Indicators

Influenza vaccine indicator status:

Automatic indication using seasonal factors The Influenza season is defined as August 1 - May 31.

Force patient to be indicated

Force patient to be contraindicated

Pneumococcal - PPV Indicator status:

PPV NOT indicated If indicated, patients under 10 years of age will be forecast three years from last Pneumococcal vaccine. Over ten years of age, the forecast interval from their last Pneumococcal vaccine is five years.

PPV indicated

Synagis® (palivizumab) indicator status:

Synagis NOT indicated If indicated, during the RSV season (October 1 - March 31) Synagis will be forecast each month from the date of last injection.

Synagis indicated

Synagis is a registered trademark of MedImmune, Inc.

Vaccine Status Recall


This report will produce a list of patients due for a Synagis injection by a selected recall date. Any patient over two-years-old will no longer appear on the report even if the indicator is still set.

1. Navigate to the Demographic Analysis/Recall report: **Practice Management tab > Demographic Analysis/Recall.**
2. Create the **Patient Demographics list**
3. Click the **Vaccine Status Recall** tab.
4. Select the **Behind or Due** radio button.
5. Click the Vaccine group drop-down and select **Synagis**.
6. Click the **Recall date** dropdown and select a date from the calendar.
7. Click **Search**.

Synagis Indicated Report

This report will produce a list of all patients that have the Synagis indicator selected in their Immunization record. The report can be used to view patients due for a Synagis injection and also display patients where the indicator should be removed. You must have the proper user permission to access OP Reports to run this report.

1. Navigate to OP Reports: **Tools tab > OP Reports.**
2. Login to OP Reports using your OP login.
3. Click the **Administration** folder in the All Folders pane.
4. Click the **Medical** folder and double click to run the **Synagis Indicated Report**
5. *(If necessary)* Click the **Additional** folder if Synagis indicated Report is not listed in the Medical folder.




Version 14.10

Overview

If the practice administers Synagis injections, it may be necessary to produce a report of patients for follow up that have the indicator set. Below are listed several reports that are available. Your practice workflow determines the report that will be used.

Note: For the below reports to produce a patient list, you must set the **Synagis indicated** status on the Indicators tab of the Immunization record.



Forecast | Complete List | Indicators | Notes | Forecast Key

Status Indicators by Antigen | Influenza / PPV / Synagis Indicators

Influenza vaccine indicator status:

Automatic indication using seasonal factors The influenza season is defined as August 1 - May 31.

Force patient to be indicated

Force patient to be contraindicated

Pneumococcal - PPV Indicator status:

PPV NOT indicated If indicated, patients under 10 years of age will be forecast three years from last Pneumococcal vaccine. Over ten years of age, the forecast interval from their last Pneumococcal vaccine is five years.

PPV indicated

Synagis® (palivizumab) indicator status:

Synagis NOT indicated


Synagis indicated If indicated, during the RSV season (October 1 - March 31) Synagis will be forecast each month from the date of last injection.

Synagis is a registered trademark of MedImmune, Inc.

Vaccine Status Recall

This report will produce a list of patients due for a Synagis injection by a selected recall date. Any patient over two-years old will no longer appear on the report even if the indicator is still set.

1. Click the **Reports** button in the main menu and select **Demographic Analysis/Recall**.
2. Create the Patient Demographics list. Click [here](#) for detail information on how to create a patient list.
3. Click the **Vaccine Status Recall** tab.
4. Select the **Behind or Due** radio button.
5. Click the Vaccine group dropdown and select **RSV-Mab**.



Patient Demographics | Clinical Recall | Care Plan Recall | **Vaccine Status Recall** | Vaccines Given

Forecast status:

Any Status Behind or Due Vaccine group: RSV-Mab

Behind Due Now / Future Recall date: 8/1/2018

Search Imm Letters

6. Click the **Recall date** dropdown and select a date from the calendar.
7. Click **Search**.

Synagis Indicated Report

This report will produce a list of all patients that have the Synagis indicator selected in their Immunization record. The report can be used to view patients due for a Synagis injection and also display patients where the indicator should be removed. You must have the proper user permission to access OP Reports to run this report.

1. Click the **Tools** button on the main menu and select **OP Reports**.
2. Login to OP Reports using your OP login.
3. Click the **Administration** folder in the All Folders pane.
4. Click the **Medical** folder and double click to run the **Synagis Indicated Report**
5. (If necessary) Click the **Additional** folder if Synagis indicated Report is not listed in the Medical folder.

