

Important Content Update Message

We are currently updating the OP Help Center content for OP 20. We appreciate your patience as we continue to make these updates. To locate the version of your software, navigate to: **Help tab > About**.

Age and Dosage Warning Messages for DTaP+IPV

Last Modified on 08/19/2020 12:53 pm EDT

Version 20.6

The Vaccine Logic in OP has been updated to create separate warning messages for DTap+IPV vaccine. The warning messages will display specific to an age or dosage inconsistency. Below you will find the messages that display. As a reminder, always check the brand when reviewing the below messages.

Warning ×	Warning X	Warning X
DBP-PVJ not indicated at this time. Status: Alert. DEP-PV9 should be administered only when: - aptient is between 4 and 6 years old (patient is 1 years 11 months old) - aptient has had a oprevious doses of DEP-PV - aptient has had 3 or more doses of DEP-PV (patient has had 3	(DT4P-IPV) not indicated at this time. Status: Alett. Ture: Ture: DT4P-IPV should be administered only when: - patient is between 4 and 6 years old (patient is 15 years 5 months 3 days old) - patient has had no previous doses of DT4P-IPV (patient has had 3 or more doses of IPV (patient has had 3 or more doses of IPV (pati	DTBP-IPV] not indicated at this time. Status: Alet. Status: Alet. Diab-IPV should be administered only when: - patient his head 4 does of DTBP or DTBP- (patient has head 3 does of OTBP or DTBP- (patient has head 3 does of OTBP Por DTBP- (patient has head 3 does of OTBP Por DTBP- patient has head 3 does of OTBP.IPV. - patient has head 3 does of OTBP.IPV. - both DTBP and IPV are Due or Behind. (DTBP: behind IPV: Due) n//e; Continue anyway?
Yes No		Yes No

